Aggregation For Dummies (Macroeconomists)

Consider an economy populated by a set of agents distributed uniformly along the unit interval with a total population mass of 1. That is, for $i \in [0, 1]$ the probability distribution function is f(i) = 1 and f(i) = 0 elsewere; the CDF on the [0, 1] interval is therefore F(i) = i, implying an aggregate population mass of F(1) = 1.

Agent i's value of variable • at date t is $\bullet_{t,i}$. Thus aggregate consumption is

$$C_t = \int_0^1 c_{t,i} f(i) di,$$

=
$$\int_0^1 c_{t,i} di,$$
 (1)

and a similar notation applies to other variables.

Since the aggregate population is normalized to 1, capital letters refer not only to aggregate variables but also to per capita variables, since per-capita consumption is aggregate consumption divided by aggregate population:

$$\left(\frac{\int_0^1 c_{t,i} f(i) di}{\int_0^1 f(i) di}\right) = \left(\frac{C_t}{1}\right) = C_t.$$
(2)

Each individual agent is infinitesimally small, and can therefore neglect the effects of its own actions on aggregates.

1 Blanchard Lives

For many purposes the assumption that economic agents live forever is useful; but for other purposes it is necessary to be able to analyze agents with finite horizons. Blanchard (1985) introduced a tractable framework that permits analysis of many of the key issues posed by finite lifetimes.

The key assumption is that the probability of death is independent of the agent's age. (This is similar to the Calvo (1983) assumption that the probability that a firm will change its prices is independent of the time elapsed since the last price change).

The most convenient formulation of the model is one in which the number of dying individuals is always equal to the number of newborn individuals, so that the population remains constant.

1.1 Discrete Time

As above, suppose that the population alive at time t is arranged on the unit interval. The probability of death is d (and the probability of not dying is $\mathcal{D} = 1 - d$). Then for a person living at any location $i \in [0, 1]$, expected remaining lifetime including the current period will be

$$1 + \overbrace{(1-\mathsf{D})}^{\not D} + (1-\mathsf{D})^2 + \ldots = \left(\frac{1}{1-(1-\mathsf{D})}\right) = 1/\mathsf{D}.$$
(3)

If a new cohort of size D has been born each period since the beginning of time, the total population will be given by the size of a new cohort D multiplied by the expected lifetime D^{-1} :

$$\mathsf{D}\mathsf{D}^{-1} = 1,\tag{4}$$

so that the mass of the aggregate population is constant at 1, as above.

1.2 Continuous Time

Blanchard's original treatment was in continuous time, with a constant rate of death d, so that the probability of remaining alive (not dead) after t periods for a consumer born in period 0 is¹

$$\aleph_t = e^{-\mathsf{d}t} \tag{5}$$

so that the expected life span is

$$\int_0^\infty e^{-\mathsf{d}\tau} d\tau = 1/\mathsf{d} \tag{6}$$

and if the flow arrival rate of new population is d (that is, at each instant a flow of new population arrives at rate d) then again the population mass is constant at

$$\mathsf{d}/\mathsf{d} = 1. \tag{7}$$

1.3 Population Growth

Now suppose that the population in the discrete-time model is growing by a factor $\Xi = (1 + \xi)$ from period to period; if the number of newborns in period 0 was 1, then the number of newborns in period t is given by

$$\Xi^t$$
. (8)

In this framework we want to keep track of the *relative* population of each cohort compared to the size of the newborn cohort. At age z, the cohort that was born in period 0 will be of relative size

$$(\not\!\!D/\Xi)^z \tag{9}$$

¹The number of people who will die during the first period in the continuous-time model is measured by the difference between an initial population of size 1 and the size of the population remaining alive after one period, $1 - \aleph_1 = 1 - e^{-d}$. [ExpEps]in MathFacts implies that $e^{-d} \approx 1 - d$; therefore the proportion who have died will be approximately 1 - (1 - d) = d. Hence for small death rates, in order for the same population to survive for one period in a continuous-time model with death rate d and the discrete-time model with death rate D, one would need $d \approx D$.

The total relative populations will be

$$1 + (\mathcal{D}/\Xi) + (\mathcal{D}/\Xi)^2 + \dots = \left(\frac{1}{1 - \mathcal{D}/\Xi}\right)$$
(10)

so that if in period 0 the population was of size $(1 - \not{D}/\Xi)$ then the sizes of the relative populations will add up to one even as the absolute population grows by the factor Ξ .

References

- BLANCHARD, OLIVIER J. (1985): "Debt, Deficits, and Finite Horizons," Journal of Political Economy, 93(2), 223–247.
- CALVO, GUILLERMO A. (1983): "Staggered Contracts in a Utility-Maximizing Framework," *Journal of Monetary Economics*, 12(3), 383–98.